Fig. 1. Landings of black coral in Hawaii. Data are pooled into seven year bins to meet confidentiality requirements for catch reporting. There is also a significant delay between harvest and reporting. Fishermen also commonly collect coral and retain this for several years and only report it after they sell the coral. Landings in the last 7 years comprised 58% of the total catch since 1985. Also, the average annual catch reported for the period 1999-2005 more than doubled the catch for the seven year period prior. Data and figure from Parrish, 2006.




Table 1. Taxa of black corals reported in CITES between 1988-2003. The dominant corals are shown in bold. Several additional taxa of black coral are reported to be in trade at low levels, but they have not been listed in the CITES database. These include the following genera: *Hillopathes Parantipathes, Taxipathes* and *Tropidopathes*

TAXA	SOURCE	VOLUME (example years)
Antipatharia spp.	All	50,000, 20 kg (2001)
Antipathes spp.	All	7000 (2001)
Antipathes crispa	US	100 kg (2002)
Antipathes grandis	All	
**Antipathes densa	All	67,000, 19 kg (2001)
Antipathes japonica	Taiwan	63,000/138,000/43,000 (1988,1992, 1996)
Antipathes columnaris	New Zealand	6 (2003)
Antipathes plantagenista	Cuba	3 (1996)
Antipathes abies	Philippines	42,700 (1996)
Antipathes dichotoma (A. cf. curvata)	Papua New Guinea Taiwan, Philippines	10,000 (1986)
Antipathes hirta	ТТ	2 (2002)
Aphanipathes spp.	U.S., Mexico	1,600 live (1995)
Bathypathes spp.	Philippines	1554 (1986) 6 (2001)
Bathypathes lyra	Russia	18 live (1997)
Bathypathes scoparia	U.S.	5 (1997)
*Cirrhipathes anguina	All	65,000 (2001)
Cirrhipathes spiralis	Papua New Guinea Philippines	270 (1987-89) 6,000 (1986)
Cladopathes spp.	Indonesia	16 live (1995); 100 (1986); 2 (2002)
Hexapathes spp.	Taiwan	1 (1996), 2000 (1987) 3 (2002)
Leiopathes spp.	Dominican Republic, Haiti, Thailand	4 (1996; 12 live (1993);152 live (1988); 3 (2001)
Myriopathes japonica	Taiwan	8000 (2003)
Parantapathes	СА	2 (2002)
Sibopathes macrospina	Australia, Fiji	1 (2002), 25 live (1995)
Stichopathes gracilis	U.S.	500 live (1996)
Stichopathes regularis	Mexico	2 live (1997).
Stichopathes longispina	Fiji	1 live (1997); 1 kg (2001)
Schizopathes spp.	СА	3 (1990) 6(2002)




Fig. 2. Total volume of imports of black coral between 1983-2003. All data are from CITES, and include only reports of black coral by piece. Additional trade in black coral is also reported by weight, with hundreds to thousands of kg reported each year.